

AD-A268 663



TAGN PAGE

Form Approved
OMB No. 0704-0188

to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering the collection of information. Send comments regarding this burden estimate or any other aspect of this form, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Avenue, Washington, DC 20540.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 1993		3. REPORT TYPE AND DATES COVERED Reprint	
4. TITLE AND SUBTITLE (see title on reprint)				5. FUNDING NUMBERS PE: NWED QAXM WU: 00145	
6. AUTHOR(S) Miller et al.					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Armed Forces Radiobiology Research Institute 8901 Wisconsin Ave. Bethesda, MD 20889-5603				8. PERFORMING ORGANIZATION REPORT NUMBER SR93-15	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Defense Nuclear Agency 6801 Telegraph Road Alexandria, VA 22310-3398				10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited.				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) <div style="text-align: center;">DTIC ELECTE AUG 30 1993 S A D</div> <div style="text-align: center;">93-19550</div> <div style="text-align: center;">93 8 23 043</div>					
14. SUBJECT TERMS				15. NUMBER OF PAGES 7	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED		18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED		19. SECURITY CLASSIFICATION OF ABSTRACT	
				20. LIMITATION OF ABSTRACT	

SECURITY CLASSIFICATION OF THIS PAGE

CLASSIFIED BY:

DECLASSIFY ON:

SECURITY CLASSIFICATION OF THIS PAGE

FREE-RADICAL YIELDS IN PROTON IRRADIATION OF ORIENTED DNA: RELATIONSHIP TO ENERGY TRANSFER ALONG DNA CHAINS

J.H. Miller*, D.L. Frasco*, M. Ye*, C.E. Swenberg**, L.S. Myers, Jr., and
A. Rupprecht***

*Pacific Northwest Laboratory, Richland, WA 99352

**Armed Forces Radiobiology Research Institute, Bethesda, MD 20814

***University of Stockholm, S-106 91 Stockholm, Sweden

ABSTRACT

Spatial patterns of energy deposition on the nanometer scale are currently believed to be a major factor in determining the biological effectiveness of ionizing radiation. If the most common precursors of biologically significant lesions are clusters of ionization in or near DNA, then intramolecular energy and charge transfer along DNA chains could be very important in lesion development. This paper describes investigations of these phenomena through model calculations and measurements of radical yields in oriented DNA exposed to proton irradiation.

INTRODUCTION

The idea that the critical lesions in radiation biology result from clustering of ionizations on a nanometer scale can be traced to the early work of Lea (1947), Howard-Flanders (1958), and Barendsen (1964). More recent support for this model of the biological effects of ionizing radiation comes from the analysis of cell killing by radiations with different linear-energy-transfer (LET) based on computer simulations of track structure in water (Goodhead, 1987). Although these biophysical models ignore the complexity of the cellular medium and the macromolecular structures that regulate its function, their basic conclusion can be rationalized by the high scavenging capacity of the chemical environment of DNA (Roots and Okada, 1975) and the efficiency of repair of minor perturbations of DNA structure (Doetsch and Cunningham, 1990).

From arguments of this type, one concludes that the most common precursor of cytotoxic and mutagenic effects from radiation exposure is a cluster of ionization in or, at least, very near to the DNA molecule. The involvement of macromolecules in the early stages of lesion production opens the possibility of intramolecular energy and charge transfer following excitation or ionization by the radiation field. These processes acting in the presence of traps for energy and charge provide a mechanism for concentrating energy deposited in macromolecular systems that is independent of stochastic processes in the slowing down of charged particles. Conversely, energy and charge transfer along DNA chains may dissipate clusters of excitation and ionization before biologically significant lesions are formed. In general, the existence of energy and charge migration in macromolecules tends to decouple lesion production from the stochastics of energy deposition

just as ordinary diffusion tends to decouple radiation chemistry on a long time scale from track effects in the radiolysis of homogeneous solutions of small molecules.

Recent findings at several laboratories have raised questions about the assumption that radiation-induced DNA damage remains localized on the nanometer scale during lesion formation. Observations by Arroyo et al. (1986) that the yield of neutron-induced free radicals in oriented DNA fibers was dependent on the orientation of the sample relative to the neutron flux were attributed to energy transfer between stacked DNA bases. Al-Kazwini et al. (1990) presented evidence that electrons can move along DNA chains for distances up to about 100 base pairs. Data obtained by van Lith et al. (1986) on microwave conductivity in pulsed radiolysis suggest that excess electrons in frozen DNA solutions can migrate for distances of the order of 100 nm in the structured water layers around macromolecular chains.

Simultaneous with these experimental results, there has been a renewed interest in nonlinear modes of vibrational excitation in DNA called solitons (Baverstock and Cundall, 1988). Several models for solitary waves in DNA have been proposed (Englander et al., 1980; Yomosa, 1984; Takeno and Homma, 1987; Zhang, 1987; Muto et al., 1988). In Yomosa's model, about 0.4 eV of vibrational energy is transported along DNA at a rate of about 100 nm/ns as a disruption of hydrogen bonds between complementary bases. The broad spectrum of excitation associated with the absorption of energy from ionizing radiation in biological systems (Bednár, 1985) should provide ample opportunity to overcome the energy threshold required to initiate nonlinear phenomena like solitons (van Zandt, 1989).

The first section of this paper discusses models for the effects of energy and charge transfer on free-radical yields in oriented DNA samples exposed to direct proton-beam irradiation. Experiments designed to detect an orientation dependence of radical yields from proton irradiation of oriented DNA are described in the second section. Our conclusions are presented in the final section.

MODELING RADICAL YIELDS IN PROTON IRRADIATION OF ORIENTED DNA

We have investigated (Miller et al., 1988; Miller and Swenberg, 1990) mechanisms for the observation by Arroyo et al. (1986) that DNA base radicals induced by neutrons are dependent on the orientation of DNA fibers relative to the neutron flux. Monte Carlo codes developed by Wilson and Paretzke (1981) and scoring algorithms contributed by Charlton (1985) were used to model energy deposition in oriented DNA under direct proton-beam irradiation. Table I summarizes our results for a 1 MeV proton flux incident on an oriented DNA sample either parallel or perpendicular to the fiber direction. Although the average amount of energy deposited in the parallel case is 5 times greater than in the perpendicular case, the average separation between the excitations or ionizations in the same DNA chain is about 100 times greater in the parallel case. The pattern of deposition in the parallel case is also more sensitive to uncertainty in the fiber orientation relative to the proton beam.

If the pattern of energy deposition events along a DNA chain in the parallel case is as diffuse as these model calculations suggests, then only long-range modes of intramolecular energy or charge transfer could couple the excitations and ionization in the molecule in ways

Table 1. Energy Deposition in Oriented DNA by 1 MeV Protons

Orientation	Energy Deposited (eV)	Event Separation (nm)
0°	293	224
0° ± 10°	150	47
90°	62	2
90° ± 10°	62	2




Figure 1. Decay Modes of Energy Absorbed in Solid DNA Samples

that would make the yield of free radicals orientation dependent. For example, recent work by Georghiou (1990) indicates that singlet excitations of bases in calf thymus DNA at room temperature move only 1 or 2 base pairs before they are irreversible trapped. Clearly, this type of excitation has a low probability of interacting with other energy deposition events in same DNA chain and should have equivalent effects in both irradiation geometries.

Energy absorbed from ionizing radiation in oriented DNA fibers should produce modes of excitation that are considerably more mobile than singlet excitons (Al-Kazwini et al., 1990; van Lith et al., 1986; Yomosa, 1984). Figure 1 illustrates a mechanism by which solitons might influence free-radical yields when DNA at 77° K is exposed to a proton beam parallel to the molecular orientation. The interaction of DNA with protons and secondary electrons produces highly excited electronic states that decay primarily through formation of electron-hole pairs. Usually this decay is accompanied by some conversion of electronic to vibrational energy. Sufficiently large vibrational excitations in DNA can be self cohering (van Zandt, 1989). Recent work by Bernhard (1989) suggests that excess electrons are trapped on cytosine by reversible proton transfer from guanine. If many super-excited states are produced in the same DNA chain by a proton flux that is parallel to the molecular orientation, then the probability of interaction between trapped electrons and solitons increases. This interaction may induce electron transfer to thymine where $TH\cdot$ is formed by irreversible protonation at C6.




Figure 2. Schematic of Electron-hole Recombination in Oriented DNA Exposed to a Proton Flux that is Parallel to the DNA Fibers.

The model illustrated in figure 1 neglects the mobility of electrons ejected in the decay of super-excited states by ion-pair formation. The high mobility and long lifetime of excess electrons in the hydration layers of DNA (van Lith et al., 1986) could also contribute to orientation effects in DNA damage by proton irradiation. This mechanism is illustrated in figure 2, where the dashed lines represent trajectories of ejected electrons that move primarily in hydration layers around DNA but occasionally are scattered between DNA chains. The squares represent fiber defects that mainly determine the mean free path of excess electrons when the sample is exposed to low LET radiation or protons perpendicular to the fiber orientation. However, when proton tracks are parallel to the fiber direction, many positive ions lie in the high-mobility path of excess electrons. This increases the probability of electron-hole recombination and could reduce the yield of primary radical anions and cation in the parallel case.

EXPERIMENTS WITH PROTON IRRADIATION OF ORIENTED DNA

Preparation of oriented DNA samples (Rupprecht, 1966) to look for effects like those illustrated in figures 1 and 2 involves wet spinning of high molecular weight DNA into fibers that are wound on a spool to form a thin sheet of oriented DNA. Samples for perpendicular




Figure 3. EPR Spectra of Oriented DNA Exposed to γ -rays (A) and Protons Perpendicular (B) or Parallel (C) to the DNA Fibers.

irradiation were made by pressing together a sufficient number of sheets to give a thickness greater than the range of a 4 MeV proton, which is about 0.5 mm. Samples for parallel irradiation were sliced from a block of oriented DNA that had a thickness slightly less than the 3 mm inside diameter of quartz tubes used to transfer irradiated samples to the electron paramagnetic resonance (EPR) spectrometer. All samples were approximately 1 cm long and weighed about 15 mg.

For irradiation, the samples were placed on a copper block in contact with a reservoir of liquid nitrogen and held in place by a thin polyester film. After cooling to 77° K, the sample was placed in a vacuum chamber attached to the beam line of the accelerator. Samples were exposed to graded doses of 4 MeV protons in the range of 20 to 60 kGy. The dose rate of 2.5 kGy/min was less than the value of 10 kGy/min recommended by Henriksen and Snipes (1970) to avoid sample heating. After irradiation, the samples were removed from the vacuum chamber and transferred to a precooled EPR tube. During this transfer, the sample lost contact with liquid-nitrogen cooled surfaces for less than one second as it fell through a funnel into the EPR tube. Several samples were exposed to γ -rays for comparison with published data (Gräslund et al., 1971) and the results of proton irradiation. In this case the sample could be sealed into an EPR tube before irradiation due to the penetrating power of the radiation.

The EPR spectra shown in figures 3A-3C were observed after 4 kGy of γ -rays, 56 kGy of protons incident perpendicular to the fiber orientation, and 48 kGy of protons parallel to the DNA chains, respectively. All three spectra are composed of a central line that we associate with primary radical anion and cation species with varying amounts of modulation in the wings resulting from TH^\cdot production. The greater amount of TH^\cdot observed with proton irradiation is not likely to be due to sample warming during the transfer from the proton beam line to the EPR spectrometer because the irradiated samples were kept in contact with liquid nitrogen cooled surfaces and the magnitude of the central line relative to the structure in the wings did not change as a function of proton dose. The greater proportion of TH^\cdot with proton irradiation may be due to the higher dose required to detect radicals since the total radical yield per unit of dose was more than an order of magnitude lower for protons than for γ -rays; however, the shape of the EPR spectrum did not change significantly with proton dose in the range investigated. The recommendation of Henriksen and Snipes (1970) regarding the dose rate was based on their experience with 6.5 MeV electrons which may not apply to proton irradiation; hence, the higher yield of TH^\cdot that we observed with protons may have resulted from sample heating due to a dose rate that was too large.

Unlike the results reported for neutrons (Arroyo et al., 1986), EPR spectra of radicals produced by direct proton irradiation of oriented DNA in parallel and perpendicular geometries were not significantly different. Figure 4 shows that, within experimental error, total radical yields were also independent of the orientation of DNA fibers relative to the proton flux. To obtain these results, differential EPR spectra were recorded digitally and




Figure 4. Total Radical Yields in Parallel (●) and Perpendicular (□) Proton Irradiation of Oriented DNA at 77° K.

double integrated to give the area under the absorption lines. This area was converted to number of spins by comparison with a standard sample of 2,2 diphenyl-1-picrylhydrazyl (DPPH) dissolved in paraffin.

CONCLUSIONS

We did not find any evidence for long-range energy or charge transfer in DNA from experiments in which oriented DNA was exposed to direct proton-beam irradiation. This may be due to the high doses and dose rates used in our experiments. The dose and dose rate could be reduced by redesigning the sample holder and transfer system to avoid the limitations on sample size imposed by the present system. Experiments with larger samples, higher proton energies for greater penetration, and improved EPR detection sensitivity might reveal orientation effects that are not present in our data due to sample heating or other processes that destroy free radicals at high exposure levels (Bernhard, 1981).

ACKNOWLEDGEMENTS

The authors gratefully acknowledge support for this work by the Office of Health and Environmental Research of the U.S. Department of Energy under contract DE-AC06-76RLO 1830, the Armed Forces Radiobiology Research Institute, and the Swedish Medical Science Research Council.

REFERENCES

- Arroyo, C.M., A.J. Carmichael, C.E. Swenberg, and L.S. Myers, Jr., 1986, "Neutron-induced free radicals in oriented DNA." *Int. J. Radiat. Biol.* 50: 789-793.
- Al-Kazwini, A.T., P. O'Neill, G.E. Adams, and E.M. Fielden, 1990, "Radiation-induced energy migration within solid DNA: The role of misonidazole as an electron trap." *Radiat. Res.* 121: 149-153.
- Barendsen, G.W., 1964, "Impairment of the proliferative capacity of human cells in culture by α -particles with differing linear-energy transfer." *Int. J. Radiat. Biol.* 8: 453-466.
- Baverstock, K.F. and R.B. Cundall, 1988, "Long range energy transfer in DNA." *Radiat. Phys. Chem.* 32: 553-556.
- Bednár, J., 1985, "Electronic excitations in condensed biological matter." *Int. J. Radiat. Biol.* 48: 147-166.
- Bernhard, W.A., 1981, "Solid-state radiation chemistry of DNA: The bases." *Adv. Radiat. Biol.* 9: 199-280.
- Bernhard, W.A., 1989, "Sites of electron trapping in DNA as determined by ESR of one-electron-reduced oligonucleotides." *J. Phys. Chem.* 93: 2187-2189.
- Charlton, D.E., D.T. Goodhead, W.E. Wilson, and H.G. Paretzke, 1985, "The deposition of energy in small cylindrical targets by high LET radiations." *Radiat. Prot. Dosim.* 13: 123-125.
- Doetsch, P.W. and R.P. Cunningham, 1990, "The enzymology of apurinic/apyrimidinic endonucleases." *Mutat. Res.* 236: 173-201.
- Englander, S.W., N.R. Kallenbach, A.J. Heeger, J.A. Krumhansl, and S. Litwin, 1980, "Nature of the open state in long polynucleotide double helices: Possibility of soliton excitation." *Proc. Natl. Acad. Sci.* 77: 7222-7226.
- Georgiou, S., S. Zhu, R. Weidner, C.-R. Huang and G. Ge, 1990, "Singlet-singlet energy transfer along the helix of a double-stranded nucleic acid at room temperature" *J. Biomol. Struct. Dyn.* 8: 657-674.
- Goodhead, D.T., 1987, "Physical basis for biological effect." *Nuclear and Atomic Data for Radiotherapy and Related Radiobiology* (International Atomic Energy Agency, Vienna), pp. 37-53.

Gräslund, A., A. Ehrenberg, A. Rupprecht, and G. Ström, 1971, "Ionic base radicals in γ -irradiated DNA." *Biochim. Biophys. Acta* 254: 172-186.

Henriksen, T. and W. Snipes, 1970, "Radiation-induced radicals in thymine: ESR studies of single crystals." *Radiat. Res.* 42: 255-269.

Howard-Flanders, P., 1958, "Physical and chemical mechanisms in the injury of cells by ionizing radiation." *Adv. Biol. Med. Phys.* 6: 553-603.

Lea, D.E., 1947, "The induction of chromosome structural changes by radiation: detailed quantitative interpretation." *British Journal of Radiology, Supplement* 1: 75-83.

Miller, J.H., W.E. Wilson, C.E. Swenberg, L.S. Myers, Jr. and D.E. Charlton, 1988, "Stochastic model of free radical yields in oriented DNA exposed to densely ionizing radiation at 77K." *Int. J. Radiat. Biol.* 53: 901-907.

Miller, J.H. and C.E. Swenberg, 1990, "Free-radical yields in DNA exposed to ionizing radiation: Role of energy and charge transfer." *Can. J. Phys.* 68: 962-966.

Muto, V., J. Halding, P.L. Christiansen and A.C. Scott, 1988, "Solitons in DNA." *J. Biomol. Struct. Dyn.* 4: 873-894.

Roots, R. and S. Okada, 1975, "Estimation of life times and diffusion distances of radicals in x-ray-induced DNA strand breaks or killing of mammalian cells." *Radiat. Res.* 64: 306-320.

Rupprecht, A., 1966, "Preparation of oriented DNA by wet spinning." *Acta Chem. Scand.* 20: 494-504.

Takeno, S. and S. Homma, 1987, "Kinks and breathers associated with collective sugar puckering in DNA." *Progress of Theoretical Physics* 77: 548-562.

van Lith, D., J.M. Warman, M.P. de Haas, and A. Hummel, 1986, "Electron migration in hydrated DNA and collagen at low temperatures." *J. Chem. Soc., Faraday Trans. 1* 82: 2933-2943.

van Zandt, L.L., 1989, "DNA solitons with realistic parameter values." *Phys. Rev. A* 40: 6134-6137.

Wilson, W.E. and H.G. Paretzke, 1981, "Calculations of distribution of energy imparted and ionization by fast protons in nanometer sites." *Radiat. Res.* 87: 521-537.

Yomosa S., 1984, "Solitary excitations in deoxyribonucleic acid (DNA) double helices." *Phys. Rev. A* 30: 474-480.

Zhang, C.-T., 1987, "Soliton excitations in deoxyribonucleic acid (DNA) double helices." *Phys. Rev. A* 35: 886-891.

Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	20

DTIC QUALITY INSPECTED 3

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

() Where copyrighted material is quoted, permission has been obtained to use such material.

() Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

(X) ^{IMK} Citations of commercial organizations and trade names in this report do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations.

() In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

() For the protection of human subjects, the investigator(s) have adhered to policies of applicable Federal Law 45 CFR 46.

() In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

Gedra M. Kowal August 14, 1993
PI Signature Date